

## Isolation and purification of hirudin from local endangered leech species

Anwar Ismail, A., Zulhisyam, A.K. and Ibrahim Che Omar

Faculty of Agro Industry and Natural Resources  
Universiti Malaysia Kelantan, Pengkalan Chepa, 16100 Kota Bharu  
Tel: +6097717234; Fax: +6097717232  
E-mail: anwar@umk.edu.my

### ABSTRACT

It has long been known that local leech (*Hirudinea* sp) contain a substance with anticoagulant properties and the isolation of this was first made in 1984 (Weinberg (1994). In this study the purification of the blood-thinning agent hirudin, protein profiling and quantification from different parts of leech body are reported. The result shows the sample from leech head gave the highest result in protein concentration with a value of 69.33 µg/ml compared with the sample from the whole body and headless leech which was 36.67 µg/ml and 26.67 µg/ml, respectively. Sample from leech head was subsequently used for purification by gel permeation chromatography on Sephadex G-75. Fractions which contain significant amounts of protein with absorbance value of more than 0.2 at 595 nm were tested for their anti-coagulant activity on rabbit blood. The result shows that Peak III (fraction number = 23, optical density = 0.256, protein concentration = 82.00 µg/ml) took the longest time for blood to clot (123.33 min) and did not contract but 380 minutes later when compared with Peaks II (fraction number = 21, optical density = 0.293, protein concentration = 94.33 µg/ml) and V (fraction number = 36, optical density = 0.214, protein concentration = 68.00 µg/ml ) where clotting was relatively earlier and occurred within 13.33 and 23.33 minutes and showed contraction much sooner at 24.33 and 41.67 minutes later, respectively. Crude protein and eluted fractions which contain significant amounts of protein with the longest time of blood to clot were subjected to SDS-PAGE analysis. The result shows that the molecular weight of hirudin was about 11.36 kDa. From the purification determined by gel permeation chromatographic technique the profiles of leech crude extract from different parts of body indicated that the anti-coagulant agent or hirudin is only found in substantial quantities in the head (salivary glands) of leeches.

*Keywords:* *Hirudinea* sp., *hirudin*, *gel permeation technique*, *protein purification and profiling*

### Introduction

Leeches (Phylum: Annelida, Class: Hirudinea) are widely distributed all over the world in a diversity of habitats, such as freshwater, seas, deserts, and oases (Gouda, 2006), where they may be important trophic links. They can be predators, vectors of parasites, as well as prey of other aquatic animals (Saywer, 1986; Keim, 1993). *Hirudinaria manillensis* is a sanguivorous, freshwater leech, with a distribution in Southeast Asia, such as in southern China, the Philippines, Thailand, Vietnam and Malaysia.

Traditionally, leeches are widely used as a model animal in toxicological, physiological, neurobiological, biochemical, histological and many other studies (Mann, 1962; Flerov and Lapkina, 1976; Lapkina and Flerov, 1979; Sawyer, 1986; Lapkina, 1992; Huguet and Molinas, 1992, 1996; Blackshaw and Nicholls, 1995; Petrauskienė, 2001). Leech therapy is one of the most important and widely practiced methods of regimental therapy used for local evacuation of morbid tumours (Ibn Sinna, 1998). The use of leeches for clinical or medicinal purposes (e.g. blood-letting) has occurred since the 5th century BC, with considerable usage in the 19th century, followed by a decline in the early 20th century (Kasperek *et al.*, 2000). Extracts (i.e. hirudins) from this species have been shown to have an important thrombolytic effect on experimental thrombosis (Tan and Liu, 2002; Huang *et al.*, 2006a, 2006b; Li *et al.*, 2006). This fact testifies to its efficiency in treating various health problems.

In recent years some leech populations have declined dramatically due to over exploitation for fishing bait and medicinal purposes and of late a further decline is observed due to environmental pollution and climate change (Sawyer, 1981; Elliot and Tullet, 1984; Wells and Coombes, 1987; Petrauskienė, 2003; Trontelj and Utevsky, 2005). However, in more recent years, the medically beneficial usage of leech has once again seen an increasing trend (Baskova *et al.*, 1983, 1992) and consequently the demand for the supply of leeches has also increased. With the rapidly expanding leech based industries the request for supply of leeches has increased greatly. The present source of obtaining leeches from their natural habitats are fast depleting primarily because of environmental pollution in many freshwater bodies a direct result of rapid industrial development in these watershed areas. To date there are various culturing methods that have been introduced but none was developed to a standard practical method suitable for breeding on a viable commercial scale. New culturing method for large scale production for local leeches is now available and which would ensure that leech farming is not a sunset industry (Zulhisyam, *et al.*, 2011).

In Malaysia, the entrepreneurs used leeches as animal for trading purposes and the stock in the form of dried powder are exported to meet the demand in countries such as Korea, Ukraine, China and Spain. On the local scene, traditional medical practitioners use the leeches for therapeutic healing treatment and is quickly gaining acceptance amongst the local folks and leech farming has thus become a thriving business. Local farmers have reported the difficulty of collecting leeches in their natural habitats of late as compared in the past and that polluted waters are the main culprit for the decline in their numbers. Several factors determine leech distribution in freshwater environments such as availability of food organisms, nature of the substrate, depth of water, presence of water currents, size and nature of the body of water, hardness and pH; dissolved oxygen, temperature of water, siltation and turbidity; and salinity (Sawyer, 1986). Polluted environment presently has upset the fine balance in the ecosystem and the leech very survival is under threat.

Hirudin is the generic name for a family of closely related homologous peptides that are all found in the cranial salivary glands of medicinal leech (Wallis, 1996). In 1984, the blood-thinning agent (anticoagulant) hirudin was isolated for the first time from leech salivary glands (Weinberg, 1994). Hirudin was extracted from the homogenized heads of medicinal leeches and enriched by precipitation procedures followed by ion-exchange chromatography and gel filtration (Badgy *et al.*, 1976; Markwart, 1957; Walsmann, 1991). Electrophoresis is a method normally used to separate and characterize a mixture of charged molecules, especially proteins and nucleic acids. SDS-PAGE is one of the common laboratory techniques in the biomedical and life sciences, and is usually used to resolve proteins in biological samples according to their size. The principle of the technique is simple, and it has been in use for many years (Laemmli, 1970).

Gel filtration (GF) separates proteins solely on the basis of differences in molecular size. Separation is achieved using a porous matrix to which the molecules, for steric reasons, have different degrees of access (smaller molecules have greater access and larger molecules less). The matrix is packed into a chromatography column, the sample mixture applied, and the separation accomplished by passing an aqueous buffer (the mobile phase) through the column. Protein molecules that are confined in the volume outside the matrix beads will be swept through the column by the mobile phase and the sample is thus eluted in decreasing order of size. Generally, the protein zones eluted are detected by an in-line UV monitor and fractions are collected for subsequent specific analysis or further preparation steps (Hagel, 1998).

## **Material and Methods**

### **Source and rearing of leeches**

Local leech used in the study was provided by PT Dynamic Consultant Co., Kota Bharu, Kelantan. The non-chlorinated freshwater in the aquarium tank placed indoor was aerated, with 50% of the water changed once every 3 days. The temperature, pH and light intensity were maintained at  $27.82 \pm 6.22$  °C,  $6.6 \pm 1.5$  and 100-150 lux, respectively. The leeches were starved for 2 weeks before starting this study.

### **Preparation of leech crude extract**

The procedure outlined by Claude (1935) was followed for the preparation of crude extract from the locally raised leeches. For this experiment, 10 g of live leeches was taken and the crops emptied of the blood by placing them in a container filled with crystals of sodium chloride for 15 minutes and then washed with distilled water. They are then cut into small pieces and grinded using a blender. The extraction was made with a volume of distilled water equal to six times the weight of the pulp. After centrifugation at 1500 rpm for 60 minutes at 4 °C, the supernatant fluid was filtered through coarse paper. Finally, the crude extract was stored at 4 °C for 24 hours. The preparation of crude extract was similarly done for leeches without head (headless) and with the leeches head comprising the first to six body segments or annuli only (Claude, 1935). The optical density (OD) value for the three parts was recorded and compared according to their protein concentrations.

### **Protein quantification**

For the preparation of the protein reagent, Coomassie Brilliant Blue G-250 (100 mg) was dissolved in 50 ml 95 % ethanol and followed with the addition of 100 ml phosphoric acid. The resulting solution was diluted with ddH<sub>2</sub>O to the final volume of one liter. Final concentration in the reagent were 0.01% (w/v) Coomassie Brilliant Blue G-250, 4.7% (w/v) ethanol, and 8.5(w/v) phosphoric acid.

As for protein assay, varying concentrations of protein solutions were prepared from the stock containing 1mg/ml BSA. The concentrations prepared were 0 (the blank), 10, 20, 30, 40, 50 and 60µl of the protein stock solution and were added to individual tubes and adjusting the volume to 100 µl with ddH<sub>2</sub>O. 1000µl of the diluted dye reagent was added to all test tubes and the content was mixed by inversion. The absorbance was measured at A<sub>595</sub> nm after 2 minutes. Standard curve was plotted for protein determination.

### **Preparation of gel-degassing**

Powdered gel weighing 7 g was added gradually to an excess amount of buffer (0.01 M sodium phosphate buffer, pH 7.0) and allowed to swell to about 84-87 ml of gel suspension and the gel supplied as slurry was suspended in a two-to fourfold excess of buffer and 95% of the gel was allowed to settle out. The remaining gel and supernatant solution was aspirated by suction. The gel was equilibrated at room temperature for 3 hours. Immediately prior to pouring the column, the gel was degassed by placing the slurry in a thick-walled Erlenmeyer flask with a side-arm. The top side was sealed with a large rubber stopper which fits over the mouth of the flask. The vacuum tubing was attached to the sidearm and connected to a vacuum line. The flask was swirled for 15 minutes and when bubbles no longer appeared, the gel was totally degassed.

### **Column assembly and packing**

The end of column was connected to a 50 ml syringe. The column was flushed with wash buffer (0.01 M sodium phosphate buffer, pH 7.0), leaving a few ml at the bottom. The column was mounted vertically on the fixture away from direct sunlight and high traffic movement. The media slurry was re-suspended and poured into the column in one continuous step. The bottom outlet of the column was opened until the gel bed reached a constant height. The column was equilibrated overnight with wash buffer until pH of gel suspension inside the column is equivalent to pH 7 (pH of wash buffer).

### **Protein separation**

As the starting material, 30 ml of crude protein extract from leech head was used with concentration of 69.33 $\mu$ g/ml. The protein determination prior to fractionation was made using a 1.5 x 30 cm BioRad Econo chromatography column filled with Sephadex G-75 which was equilibrated overnight with 0.01 M sodium phosphate buffer, pH 7.0. The protein was loaded onto the column and eluted twice the column volume (80ml) of the equilibration buffer. Fractions (3 ml per tube) were collected throughout the experiment and absorbance of proteins was measured at 595 nm by spectrophotometer (Thermo scientific). Protein concentrations were determined separately for each fraction with absorbance value of more than 0.2 at 595 nm measured by Coomassie protein assay. The procedure was repeated on samples obtained from headless leech and the whole body with the initial protein concentration of 26.67 and 36.67 $\mu$ g/ml respectively.

### **Protein activity and profiling**

Fractions which contain significant amounts of protein with absorbance value of more than 0.2 at 595 nm were tested for their anti-coagulant activity by adding 0.5 ml of the extract to 2 ml of fresh rabbit blood and the time of clotting recorded by using the drop method (Prakasam *et al.*, 2005). Crude protein and eluted fractions which contain significant amounts of protein with the longest time of blood to clot were subjected to SDS-PAGE analysis using MP3 Bio-rad minigel apparatus with 12% resolving buffer. The crude extract was mixed with 1:1 of sample buffer [13.16 % (v/v) 0.5 M Tris-HCl, pH 6.8, 26.32% (v/v) glycerol, 21.05% (w/v) SDS, 2.11% (w/v) Bromophenol blue]. The mixture was then boiled for 5 minutes. Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS-PAGE) was prepared as described by Leammli (1970). Electrophoresis was carried out in running buffer [0.3% (w/v) Tris-base, 1.44% (w/v) glycine, 0.1% (w/v) SDS; pH 8.3] at constant current of 100 V for 75 minutes. The gels were then stained with Coomassie Brilliant Blue R-250 [0.2 % (w/v) in 28% (v/v) isopropanol and 11% (v/v) acetic acid] overnight with gentle agitation at room temperature and de-stained with

solution containing 10% (v/v) acetic acid and 10% (v/v) isopropanol for 1 hour. Molecular weight was evaluated by making comparison with standard molecular weight markers (SDS-PAGE Molecular Weight Standard, Low Range, Bio-Rad).

## Results

### Protein quantification

Protein quantification was carried out by using the assay method adopted by Bradford with Bovine Serum Albumin (BSA) as a standard to determine unknown concentration of substance in particular volume. Table 1 shows the protein concentration of leech crude extract from different body parts of leeches.

Table 1. Protein concentration of extracts from different body parts

Sample	Total crude (ml)	Protein concentration ( $\mu\text{g/ml}$ )	Protein (mg)
Whole body	30	36.67	1.10
Headless leech	30	26.67	0.8
Leech head	30	69.33	2.08

The result shows the sample from leech head gave the highest protein concentration of about 69.33  $\mu\text{g/ml}$  compared with either the sample from the whole body or headless leech. Similarly for the total protein content, the sample extracted from leech head produced 2.08 mg protein far more in quantity than from either of the two sources.

### Protein profiling

Sephadex G-75 Gel Filtration chromatography technique was used to separate the protein fractions for purification of the hirudin protein. The protein profile of the fractions from the crude extract obtained from leech head is shown in Figure 1. Four distinct peaks were obtained, namely Peaks II – IV. That for the fraction at the first peak eluted during the wash was assumed as representing proteins that did not adhere to the column. The second, third, fourth and the fifth peaks represented proteins that adhered to the column very well. The result shows that Peak II gave the highest optical density (OD) reading with a value of 0.293 compared with Peaks III, IV and V with optical density values of 0.256, 0.18 and 0.214 respectively. Only peaks with OD readings greater than 0.2 were chosen for subsequent protein quantification.




Figure 1. Purification of protein obtained from leech head.

For protein quantification Bradford assays with Bovine Serum Albumin (BSA) was used as a standard to determine unknown concentration of substance in a particular volume. Table 2 shows the different protein concentration for the corresponding peaks.

Table 2. Protein concentration of crude extracts of leech head for corresponding peaks.

Peaks	Fraction Number	Optical density (O.D)	Protein concentration ( $\mu\text{g/ml}$ )
I	-	-	-
II	21	0.293	94.33
III	23	0.256	82.00
IV	-	-	-
V	36	0.214	68.00

\*Peak I: Assumed as represented proteins that did not adhere to the column

\*Peak IV: The O.D value < 0.2 not considered for determination of protein concentration

The result shows that the sample from Peak II and the fraction number 21 gave the highest protein concentration of 94.33  $\mu\text{g/ml}$  compared with the samples from the Peaks III (fraction number 23) and V (fraction number 36) which was 82.00  $\mu\text{g/ml}$  and 68.00  $\mu\text{g/ml}$ , respectively.

The protein profile of the fractions from the crude extracts obtained from whole body produced five distinct peaks as is shown in Figure 2.




Figure 2. Purification of protein obtained from the whole body of leeches.

The result shows that Peak I produced the highest optical density with a value of 0.197 (fraction number 20) compared to the other four peaks and the lowest at Peak III with a value of 0.06 (fraction number 29). As the optical density values for all peaks were below 1.5 there was no necessity to determine the protein concentration of the sample.

Similarly, the protein profile of the fractions obtained from crude extract of headless leech also produced five distinct peaks but at different fraction numbers to that profile shown by extracts obtained from whole body (Figure 3). The first peak eluted during the wash was assumed to represent proteins that did not adhere to the column. As the optical density values for all peaks were below than the required value of 0.2 there was therefore no necessity to determine the protein concentration of this extract.

For subsequent experiments only samples from crude extracts obtained from leech head was used as the source for protein since the optical density value obtained was greater than 0.2 compared with two other sources.




Figure 3: Purification of protein obtained from headless leeches.

#### Determination of hirudin protein activity

As described earlier, the sample of the crude extracts of leech head was obtained from purification by using Sephadex G-75 Gel Filtration chromatography technique and Table 3 shows the time taken for blood to clot and contract at three different peaks. The activity of protein hirudin was determined based on the time of blood to clot.

Table 3. Time taken for blood to clot and contract from samples of crude extracts of leech head obtained at three different peaks

Peaks	Clotting time (min)	Time taken for blood clot to contract (min)
II	13.33±2.89	24.33±0.58
III	123.33±15.28	380.00±17.32
V	23.33±2.89	41.67±2.89
Control 1	1±0.0	1.17±0.29
Control 2	3.33±0.58	3.33±0.58

Data in the table were means and standard deviations (mean ± S.D.)

Peak II: Fraction number = 21, O.D = 0.293, protein concentration = 94.33 µg/ml  
 Peak III: Fraction number = 23, O.D = 0.256, protein concentration = 82.00 µg/ml  
 Peak V: Fraction number = 36, O.D = 0.214, protein concentration = 68.00 µg/ml  
 Control 1: Blood rabbit only  
 Control 2: Blood rabbit treated with distilled water

The result of present study showed that the Peak III gave the longest time for blood to clot (123.33±15.28 minutes) and only contracted 380.00±17.32 minutes later when compared with Peaks II and V where the blood tested was shown to clot in a much shorter time of 13.33±2.89



and  $23.33 \pm 2.89$  minutes and contracted much sooner within  $24.33 \pm 0.58$  and  $41.67 \pm 2.89$  minutes, respectively. The control treatment with pure rabbit blood gave the lowest result when clotting occurred within  $1 \pm 0.0$  minute by which time contraction also took place.

### Protein profiling

The crude protein bands of the crude extracts of leech head on 12 % SDS-PAGE is presented in Figure 4.




Figure 4. Electrophoretic patterns (12% polyacrylamide gel) on crude extract from leech head. Lane 1-3: crude extract from leech head loaded with  $0.69 \mu\text{g}$  per well. M: SDS-PAGE Molecular Weight Standards, Low Range (BIO-RAD).

The figure shows a broad range of protein ranging from more than 97.4 kDa to less than 14.4 kDa. From the electrophoretic bands shown on the 12% polyacrylamide gel, there were 9 profile bands of unknown proteins from the leech head sample before purification was made.

The protein profile after purification was carried out is shown in Figure 5. Only one band appeared after purification and the molecular weights of the band was about 11.36 kDa. The molecular weights of the unknown protein were determined by using the curve generated by plotting the log of the molecular weight of the broad range standard vs. the relative mobility (Hames & Rickwood, 1990).




Figure 5. Electrophoretic patterns (12% polyacrylamide gel) on purification of crude extract sample of leech head by Sephadex G-75 Gel Filtration Chromatography. (Peak III: Fraction Number = 23, O.D = 0.256, protein concentration = 82.00  $\mu\text{g/ml}$ ). Lane 1: protein extract from leech head loaded with 0.82  $\mu\text{g}$  per well. Lane M: SDS-PAGE Molecular Weight Standards, Low Range (BIO-RAD).

## Discussion

In this study, new information on detection and purification of hirudin protein from local leeches are reported. Accurate determination of protein concentration is fundamental for all quantitative measurement of biochemical interactions. Before the purification process was carried out, initial studies on protein content was conducted to determine in which part of the leeches anatomy contained large concentration of protein. The protein concentrations between different body parts of local leeches were determined by Bradford assay method with Bovine Serum Albumin (BSA) which was used as a standard to determine unknown concentration of a substance in a particular volume and absorbance of proteins measured at 595 nm. This is different with the study conducted by Orevi (1992) who used the method adopted by Lowry *et al.* (1951) to determine the protein concentration from the saliva of *Hirudo medicinalis* and protein absorbance was measured at 280 nm. Generally, the Bradford method will only detect proteins with a molecular weight greater than 3-5 kDa and this particular protocol is recommended for 20-140  $\mu\text{g}$  protein (200-1400  $\mu\text{g/ml}$ ) (Rosenberg, 1949). The results of present study showed

that leech head gave the highest protein concentration of 69.333 µg/ml compared with the sample from either the whole body or headless leech (Table 4.5).

Although Walsmann (1991) on the heels of what Markwardt (1957) had achieved finally succeeded in the preparation of a nearly 50% pure hirudin sample. However, the way to reach this goal was long and stony and, as a first step required the development of appropriate chemical extraction procedures to obtain sufficient amounts of starting material and this material was the head of the leech. In the original report on some properties of hirudin, Bodong (1905) noted that 10% of a leech head contained sufficient hirudin to keep rabbit blood uncoagulated *in vitro* for more than two days. This statement was supported by Claude (1937) who argued that extracts prepared from the leech head were usually purer than those prepared from the whole body. In his study, the effects of extracts from different parts of the leech on skin permeability resulting in the greatest spread was obtained from the isolated head and gonads with the area of spread from 0.5 ml of extract with 0.25 ml India ink (as an indicator) was 91.2 and 80.7 sq. cm., respectively, compared with the extract from leech body which was only 66.9 sq. cm. According to Nowak *et al.* (2007) about 600 mg of natural pure hirudin could now be prepared from about 15,000 leech heads.

The purification process of leech crude extract was continued with the use of Sephadex G-75 gel filtration. Gel filtration chromatography is also known as molecular sieving, gel permeation and size exclusion chromatography (SEC). The primary objective of gel filtration is to achieve rapid separation of molecules based on size (Rosenberg, 1949). For the Sephadex G-75 Gel Filtration, the useful working range (molecular weight) was around 3 kDa to 70 kDa. In this study, the size of the columns used was 1.5 x 30 cm. The height of the dextran filling was about 25cm. The main advantages of the gel chromatographic technique presented in this paper over the existing techniques are that it is rapid and simple to operate. The columns can be used repeatedly over a considerable period of time. The bed volume can be maintained constant by adding freshly soaked Sephadex G-75 for any lost by leaching during the washing procedure. After the solution has penetrated the gel bed the latter was carefully covered with buffer and the elution started. The gel bed will be cracked when dry and it cannot be used again. In the present study, the first 44 ml was eluted with wash buffer and the assumptions made that the represented proteins did not adhere to the column very well. In the study conducted by Verena-Steiner *et al.*, (1990), initial purification of leech crude extract by gel permeation chromatography on Sephadex G-50 and anion-exchange chromatography on Q Sepharose fast-flow removed most contaminants and yielded a highly active extract. The result of the present study showed that purification by using Sephadex G-75 gel filtration also gave clear indication that most of the contaminants were also removed according to the large result obtained in protein quantification and SDS-PAGE analysis. Generally, the high peak does not necessarily indicate that it contained a high amount of hirudin protein. For example, as can be seen in Table 3 although the Peak II gave the highest optical density (0.293) and highest in protein concentration (94.33 µg/ml), the time taken for blood to clot when treated with leech extract was only 13.33 minutes when compared with the Peaks III and V which was 123.33 and 23.33 minutes respectively. The result clearly indicates that this particular local species not only has the hirudin protein from its extract but it also contained many other unknown proteins. In this respect, more detailed studies should be made on this local species to ascertain the molecular structure and function of some of the unknown proteins.

Generally, activity of hirudin was determined based on its inhibition to thrombin-catalyzed reactions and expressed as antithrombin units (ATU) (Grießbach *et al.*, 1985 and Spanagel *et al.*, 1991). Markwardt (1970) stated that the fast and specific reaction of hirudin with thrombin was utilized for quantitative determination of the inhibitor. He stated that one mole

of hirudin complexes with 1 mol of thrombin. This corresponds to the antithrombin activity of pure hirudin of which 1 µg inhibits about 5 µg of human thrombin. Therefore, the activity of hirudin is measured in antithrombin units (ATU), where 1 ATU is the amount of hirudin that neutralizes 1 IU of thrombin. He also stated that pure hirudin contains approximately 10,000-15,000 ATU/mg of protein, depending on the thrombin used for standardization. In this study, the protein was tested on their anticoagulant activity by contact the leech sample to the blood rabbit and time of blood clotting was recorded. We used this method because it is a very simple method and we assumed as the longest time taken for blood to clot means the highest anticoagulant agent or activity in that sample. This simple method was also used by Claud, A. (1937) in his study to determine the spreading and anticoagulant properties of the leech extract.

Electrophoretic methods have become important tools for analyzing and characterizing macromolecules. According to Weber & Osborn (1969), 0.01 mg of protein was usually applied per gel, but the amount of protein could be lowered if the gel was stained for longer period. In this study, the amount of protein loaded from leech head was 0.82 µg sample into the gel, but the gel was stained overnight. Polyacrylamide gel electrophoresis had been carried out to separate proteins according to their sizes or molecular weight and this method has been optimized to get clear banding pattern of protein profile. Optimization of the gel percentage to 12% produced the clearest banding pattern from SDS-PAGE analysis where the separation was carried out at a constant current of about 400mA starting at 100V which lasted about 90 minutes until the band's marker had reached the bottom of the gel.

To date, there are very few reports on the molecular weight of hirudin protein. In the present study, the peak that gave the longest time of blood to clot (Peak III: fraction number = 23, optical density = 0.256, protein concentration = 82.00 µg/ml) was chosen for electrophoretic analysis and the result shows that there was only one band appearing on the 12 % gel electrophoresis with the molecular weight of 11.36 kDa compared with the protein profile from unpurified extract which produced 9 bands of unknown protein. SDS-PAGE analysis (Figure 4 and Figure 5) showed that purification of leech crude extract was achieved by using Sephadex G-75 Gel as evidenced by the bands that appeared before purification compared with that after purification. Baskova (1976) determined the molecular weight of a hirudin purified by electro-focussing to be of 7.1 kDa which was close to the molecular weight determined from amino acid sequence. She also found that hirudin can dimerize in aqueous media and detected the molecular structure in a highly purified preparation with the two-fold molecular weight of about 14 kDa. In addition, Markwardt (1992) stated that the pure anticoagulant agent obtained is a carboxyhydrate-free single chain mini-protein containing 65 amino acids with a molecular weight of about 7kDa. Tan *et al.* (2002) studied the effect of geographical factors on the hirudin gene's variation of *H. manillensis*. By isolating genomic DNA and amplifying hirudin gene by the PCR method, cloning and measuring sequence as well as comparing *H. manillensis* from Guangdong Province with the hirudin gene, *H. manillensis*, from Manila, it was discovered that the geographical factors have variation effect on the hirudin gene of this leech.

In this study, the molecular weight of hirudin was found to be about 11.36 kDa and this fact is reinforced by all other information gathered where the molecular weight of hirudin in this study was similar to those of other studies and what small differences observed might be due to the different leech species used. This study is one of the few conducted to verify the presence of an anticoagulant agent, hirudin, whose molecular weight was found to be 11.36 kDa. This evidence alone may not be sufficient to provide a satisfactory explanation on the actual molecular weight of hirudin without first conducting proper molecular studies on the local leeches.




Figure 6 : Comparison of the profile of crude extracts from different body parts.

(The blue line — represented purification of protein obtained from leech head, the red line — from the whole body and the black line — from the headless leech).

Hirudin is the generic name for a family of closely related homologous peptides and usually found in the cranial salivary glands of medicinal leech, a fact confirmed by several workers (Wallis, 1996; Weinberg, 1994). In the present study, it is evident that the anti-coagulant agent (hirudin) was also from the salivary glands located in the leech head.

## Conclusion

The purification process of the crude extract of local leeches by chromatographic gel electrophoresis technique indicated the presence of the hirudin protein which is largely found in the head where the salivary glands are located. This protein when tested on pure rabbit blood demonstrated its anti-coagulant activity whose molecular weight was found to be 11.36 kDa.

## Acknowledgements

The author is grateful to the Faculty of Agro Industry and Natural Resources for providing space and facilities to carry out the study which was funded by Universiti Malaysia Kelantan under the research grant **R/SGJP/A03.00/00193A/001/2010/000031**. This research was supported by PT Dynamic Consultant Co., Kota Bharu, Kelantan and the Freshwater Fisheries Research Centre, Gelami Lemi, Jelebu, Negeri Sembilan.

## References

Bagdy, D., Barabas, E., Graf, L., Peterson, T.E., & Magnusson, S. (1976). Hirudin Methods. *Enzymol*, 45, 669-78.

- Baskova, I.P., Cherkesova, D.U., & Mossolow, V.V. (1976). Purification of hirudin by isoelectric focusing. *Biochimija*, 41, 939-941.
- Claud, A. (1937). Spreading properties of leech extracts and the formation of lymph. *Journal of Experimental Medicine*, 66. Retrieved Jan 20, 2009 from <http://www.jem.rupress.org//articles>
- G. Nowak & K. Schror (2007). Hirudin - the long and stony way from an anticoagulant peptide in the saliva of medicinal leech to a recombinant drug and beyond. *Thromb Haemostology*, 98, 116-119.
- Grießbach, U., Stürzebecher, J., & Markwardt, F. (1985). Assay of hirudin in plasma using a chromogenic thrombin substrate. *Thrombosis Research*, 37, 347-50.
- Hagel, L. (1998). Gel-filtration chromatography. In: *Current Protocols in Protein Science* (pp. 8.3.1-8.3.30). New York: John Wiley & Sons.
- Hames, B.D., & Rickwood, D. (1990). *Gel Electrophoresis of Proteins. A Practical Approach* (2<sup>nd</sup> Ed) New York: Oxford University Press. pp. 17.
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227: 380-685.
- Lowry, O.H., N.H. Rosebrough, A.L. Farr, & R.J. Randall (1951). Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193: 265-275.
- M. Orevi, M. Rigbi, E. Hy-Em, Y. Matzner, & A. Eldor (1992). A potent inhibitor of platelet activating factor from the saliva of the leech *Hirudo medicinalis*. *Prostaglandins*, 43: 483-495.
- Markwardt, F. (1992). Hirudin the promising antithrombotic. *Cardiovascular Drug Reviews*, 10 (2): 221-232.
- Markwardt, F. (1970). Hirudin as an inhibitor of thrombin. *Methods Enzymol.*, 19: 924-32.
- Markwardt, F. (1957). Die Isolierung and Chemische Charakterising des Hirudins. Hoppe Seylers Z. *Physiological Chemistry*, 308: 147-156.
- Markwardt, F. (1957). Die Bestimmung des Thrombins durch Titration mit Hirudin. *Arch Pharmazie*, 290: 280-284.
- Prakasam, L. Reddy, L. Gowri Sankar Reddy & L. Vidya Reddy (2005). *Practical Physiology: Bleeding time and clotting time*. Paras Medical Publisher, India.
- Rosenberg, Ian M. (1949). *Protein Analysis and Purification* (Benchtop Techniques). Boston: Birkhauser.
- Spanagel, M., Bichler, H., Lill, H., & Schramm, W. (1991). A fast photometric assay for the determination of hirudin. *Haemostasis*, 21: 36-40.

- Tan, E. G., Liu, X. P. (2002). Cloning and sequencing of hirudin gene of *Hirudinaria manillensis* in Guangdong, China. *Journal of Sun-Yatsen University (Medical Science)*, 23(2): 84–86.
- Verena Steiner, Rene Knecht & Markus Gruetter (1990). Isolation and purification of novel hirudins from the leech *Hirudinaria manillensis* by high-performance liquid chromatography. *Journal of Chromatography*, 530: 273-282.
- Waber, K., & Osborn, M. (1969). The realibility of molecular weight determination by dodecyl-polyacrylamide gel electrophoresis. *The Journal of Biological Chemistry*, (16): 4406-4412.
- Wallis, R. B. (1996). Hirudin from leeches to man. *Sem. Thromb. Haemost.*, 22: 185–196.
- Walsmann, P. (1991). Isolation and characterisation of hirudin from *Hirudo medicinalis*. *Sem Thromb Haemost.*, 17: 83-87.
- Weinberg, S. L. (1994). New medicine from old-hirudin and the leech. *Journal of American College of Cardiology*, 23: 544-545.