

Indigenous Plants and Tuber as Alternative Food Sources for Temiar Communities During Flood Situations

¹Mohammad Amizi Ayob, ¹Aminah Nutman, ¹Lee Lih Huey, ¹Anidah Alias, ²Shamsul Khamis, ¹NurEleenaFarihinRedzuan Hafiz Boon, ¹Tew Sue Teng and ¹Soon Jan Mei

¹Faculty of Agro Based Industry, Universiti Malaysia Kelantan, 17600 Jeli, Kelantan, Malaysia

²Institute of Bioscience, University Putra Malaysia, 43400 Serdang, Selangor, Malaysia

Address For Correspondence:

Mohammad Amizi Ayob, Faculty of Agro Based Industry, Universiti Malaysia Kelantan, 17600 Jeli Kelantan, Malaysia.
Tel: +6019-5737933; Email: soon_jan@umk.edu.my

This work is licensed under the Creative Commons Attribution International License (CC BY). <http://creativecommons.org/licenses/by/4.0/>



Open Access

Received 28 December 2015; Accepted 28 January 2016; Available online 24 February 2016

ABSTRACT

Temiar tribal people are famous for using forest resources as their food and medicinal supply. During the 2014 monsoon flood, affected villagers evacuated to nearby hills to escape the flood and rising river water. Hence, Temiar communities who used to practise the utilization of indigenous plants and tubers were forced to search for alternative food sources during the flood. Hence it is the objective of this paper to determine the types of indigenous plants and tubers used as food sources by the affected villagers. Sampling of plants consumed by the villagers were conducted at selected villagers. Voucher specimens were created and sent for identification. Fourteen types of plants and tuber were identified as alternative food sources by the Temiar communities used during the natural disaster. Plant species include paddy (*Oryza glaberrima*), tuber (*Manihot esculenta*), shrubs (*Solanum torvum*, *Clidemia hirta*), palm (*Arenga obtusifolia*), herbaceous (*Erechtites valerianifolia*, *Eryngium foetidum*, *Pentaphragma begoniifolium*), macrofungus (*Auricularia* sp.) bamboo shoot (*Dendrocalamus asper*), wild fruits (*Castanopsis megacarpa*, *Molineria capitulata* and *Rubus sumatranus*) and water extracted from stem of *Caesalpiniasappan*. Wild local plants were readily available and contributed nutrients and energy to the villagers. The identification of 14 types of food plants and tuber will provide alternative food sources for the Temiar ethnic groups particularly during lack of food availability and accessibility.

KEYWORDS: food security; natural disaster; nutrition; Orang Asli; wild plants

INTRODUCTION

The monsoon flood hit the East Coast of Malaysia in December 2014 and January 2015 and was one of the worst for the past three decades. Over 250,000 people were evacuated and there were 21 deaths. A minimum of 14 orang asli regroupment schemes were affected by the 2014 floodwaters. Accessibility to basic needs (food, medicine and relief centres) were affected as the floodwaters damaged roads and bridges near their villages (The Star, 2015) [1] (Figure 1). The Temiar communities were affected as food aid were slow to reach their villages. This led to reduced food trade (particularly for rice, salt, canned food and sugar) among the communities.

Temiar tribal people are famous for using forest resource as their food and medicine supply (Mohammad and Sharif, 2012) [2]. In Kelantan, forest plants had been used as indigenous food by the locals but are no longer practised by the younger generations. The elderly Temiar communities, village head and medicine man / woman are knowledgeable about the type of edible indigenous plants. During the flood, affected villagers evacuated to nearby hills to escape the flood and rising river water. Hence, Temiar communities who used to practise the utilization of indigenous plants and tubers were forced to search for alternative food sources during

the flood. During this period, the younger generation and men were no longer adapt in identifying edible plants and tubers. Via this flood disaster, the older generations of the Temiar tribes actively searched for alternative plants and tubers as food. The younger generations of the Temiar communities were re-introduced to the various types of edible, indigenous plants. Besides, traditional plant foods are sources of inexpensive but high quality nutrition for the local community. These plants grow wild and are readily available in the field as they do not require any cultivation (Odhav *et al.*, 2007) [3].



Fig. 1:

Medicinal plants and vegetables that grow wildy are readily available and represent inexpensive sources of high-quality nutritional food for the indigenous community. Vegetables are a rich source of vitamins and contribute to antioxidant activity in the diet [4] (Gupta and Bains 2006). Consumption of a variety of traditional plants will contribute to a diversified diet needed to meet daily micronutrient requirements. Recent studies [3, 5, 6, 7, 8] (Dias *et al.*, 2014; Mutheeswaran *et al.*, 2011; Odhav *et al.*, 2007; Toyang and Verpoorte, 2013; Wang *et al.*, 2014) had stressed on the nutritional value of traditional plants and leafy vegetables. Traditional leafy vegetables were found to contain vitamin A (99 – 1070 μg retinol equivalent), ascorbic acid (2 – 198 mg), riboflavin (0.1 – 0.6 mg) and folate (16 – 457 μg) [9] (Uusiku *et al.*, 2010). Minerals such as iron ranged from 3 – 16.2 mg/100g, zinc (0.7 – 1.0 mg/100g), magnesium, (34.5 – 141 mg/100g) and calcium (151 – 586 mg/100g) [10] (Schonfeldt and Pretorius 2011).

Tuber roots are also good sources of energy while tapioca or cassave leaves provide protein, vitamins and minerals. Tapioca roots can provide up to 150 kcal, 3.5 g protein and 3.7 g fiber /100g. Vitamins such as Vitamin A (5.0 – 35.0 μg), thiamin (0.03-0.28 mg), riboflavin (0.03-0.06 mg), niacin (0.6-1.09) and ascorbic acid (14.9-50 mg) can be found in roots [11, 12] (Charles *et al.*, 2005; Montagnac *et al.* 2009). Tapioca can be found in the wild or grown as food reserve and need not be harvested if domestic food supplies are plentiful [13] (FAO 1990). Temiar households can opt to cultivate root crops as one of the hunger mitigation strategy during flood season.

Nutritious plants and tubers can be utilised as alternative food, especially during flood seasons of due to inaccessible routes. However, in order to ensure the local communities are aware of the types of edible plants and tubers, the plants should be re-introduced to the community – particularly to the young (ages 18 – 35 years) and middle-aged adults (age 36 – 55 years). These two age groups are significant as they play a role in ensuring household food security, contribute to livelihood sustainability of the village decision-making roles. Hence it is the objective of this study to identify potential plants and tuber which can be utilised as alternative food sources during periods of food insecurity or emergency situations.

Methodology:

Identification of sites affected by 2014 floodwaters:

Affected villages (Kg. Jedip, Pos Brook; Kg. Teranek, Kuala Betis and Kg. Gob, Pos Gob) were selected as the main sites of study (Figure 1). There are a total of 2388 Temiar family households in Gua Musang including some Medriq tribes (60 households) in Pos Kuala Lah bordering Gua Musang and Kuala Krai [14] (JAKOA 2014). The villages were selected based on whether their villages were affected by the monsoon flood, accessibility (sedentary areas) and recommendations from Jabatan Kemajuan Orang Asli (JAKOA). Permission to conduct sampling at the selected villages were obtained from JAKOA and village heads.

Sampling of indigenous plants and tubers used as alternative food sources during the flood:

Field trips were conducted at the hill sites where locals collected the plant food they consumed during the flood. All plant species cited for food purposes were collected from the field. Names of local plant food and tubers, their utilized parts and preparation methods were recorded. The following table was used to record the plant information. In depth interviews with the village representatives (head of tribesmen) and medicine men / women were conducted.



Fig. 2: Selected villages affected by the flood

Herborization:

Samples were dried, and identified taxonomically based on literature. Each herbarium specimen should include important parts such as leaves, stems, flowers and fruits whenever available. For small herbaceous plants, the whole plant should be collected [15] (Semenya and Maroyi 2012). Unidentified plants were sent to Bioscience Institute, Universiti Putra Malaysia for identification and verification.

RESULTS AND DISCUSSION

Identification of indigenous plants and tubers as alternative food sources during flood:

A total of 14 plant samples were identified in all three sites. Local Temiar and Malay names were recorded. Parts used and preparation methods were documented based on the interviews with the village head of tribesman. Table 1 shows the demographic characteristics of the respondent who participated in the in-depth interview while Table 2 reveals the types of plants utilised as alternative food sources.

Some of the indigenous, edible wild plants were also found elsewhere. For example, *Auricularia auricula-judae* (jelly ear fungi) are consumed in the forests of Oaxaca, Mexico [16] (). In Malaysia, macrofungi such as *Auricularia* sp., *Clavulina* sp., *Schizophyllum commune*, *Termitomyces microcarpus* and *Termitomyces* sp. are considered edible by 5 sub-tribes: namely Semai, Temuan, Che Wong, Bateq and Jakun. Based on Lee *et al.* [17], *S. commune* was the most widely consumed fungus but most tribes prefer *Clavulina* sp.

During the flood, the affected villagers were able to access wild fruits growing in the forests. The fruits of *S. torvum* are edible and commonly found in local Malay and Thai cuisines. It is considered an important medicinal plant in tropical and subtropical countries [18]. On the other hand, there has been very little report on the consumption of *C. megacarpa*. A study conducted by Yasuda *et al.* [19] revealed that this prickly fruit is consumed by frugivorous mammals of Malaysia (e.g. leaf monkey, squirrels). Meanwhile, *Rubus sorbifolius* from the *Rosaceae* family are usually eaten raw by the Hmongs [20] and Mien [21] tribes of Thailand. In fact the leaves and stems of *R. sorbifolius* are used to treat severe cough and tuberculosis among the Mien ethnic group. In the Philippines, the Kalanguya tribe uses the extracts of *M. capitulata* as shampoo and body care [22] while the species was used for medicinal purposes in Malaysia [23].

Leaves of *E. valerianifolia* also known as gipun in Sarawak, Malaysia are used as plants for food. [24]. Alternative drinking water can be extracted from *C. sappan*, a type of heartwood which is used in Kerala, India as herbal drinking water to quench thirst, purify blood and to improve complexion [25]. A decoction of Sappan heartwood is commonly used to treat arthritis [26] and as a traditional drink or jamu material in Indonesia [27]. Bamboo shoots such as *D. asper* were found to contain high nutritive content such as carbohydrate (2.90 g/100 g), protein (25.80 g/100g dry weight), vitamins C and E (3.20 and 0.91 mg/100 mg fresh weight) [28]. In fact, *D. asper* is reported as one of Thailand's favourite edible shoots [29]. Other wild plants consumed during the flood include *Clidemia hirta*, a type of invasive perennial shrub. The fruits of *Clidemia hirta* are edible and are also consumed in certain parts of Madagascar [30]. Based on Styger *et al.* [30] studies, the existence of wild fruits represents an important supplement to the daily diet. People usually consume the fruits while traveling through the forest and are collected during shortage of food period. Sayurkelap

(*Pentaphragmabegoniifolium*) consumed in Kuala Betishas been reported as a rare and odd perennial herb that grows in the lowland rainforests of Malaysia [31]. There are limited published reports on this herb.

Table 1: Demographics of village representatives and Tok Batin (medicine men / women).

| Demographics | N = 6 |
|--|---|
| Gender | |
| Male | 5 |
| Female | 1 |
| Role in village | |
| Head of tribesman | 3 |
| Assistant to head of tribesman | 1 |
| Medicine man | 1 |
| Medicine woman | 1 |
| Others (please state): | 0 |
| Villages inundated with water | 0 |
| Yes | 2 |
| No | 0 |
| Connecting roads / bridges affected by floodwaters? | |
| Yes | 4 |
| No | 0 |
| Experience difficulty in receiving food aid during the flood? | |
| Yes | 6 |
| No | 0 |
| Searched for alternative food sources in the jungle / hills during the flood | |
| Yes | 6 |
| No | 0 |
| Total types of wild, indigenous plants consumed during the flood | |
| Below 3 types | 4 |
| 4-6 types | |
| 7-9 types | 2 |
| More than 9 types | |
| Did you consume other types of food (besides wild, indigenous plants)? | |
| Yes | 6 |
| No | 0 |
| Types of other food consumed | |
| Rice | 6 |
| Canned food | 6 |
| Crackers | 6 |
| Tea, coffee, milk or beverages | 6 |
| Others (please state): | Bread, candies, cooked dishes (at relief centres) |
| Did you hunt for animals such as squirrels, birds, wild boars or monkeys during the flood? | |
| Yes | 0 |
| No | 6 |

Meanwhile, stems / shoots of *Arengaobtusifolia* are used as traditional food of the Adi tribe in India during droughts and for preparing their local beverages [32]. Alternative food such as cassava (*M. esculenta*) and upland rice (*O. glaberrima*) are essential sources of carbohydrate and provided much needed energy for the villagers during the flood period. The starchy roots of cassava is a well-known major diet in Africa, Asia and South America [33]. Even though upland rice variety produces lower yield, it features good characteristics such as fragrance, size, shape and colour. Additionally upland rice is able to survive on hilly areas with little access to water [34]. *Eryngium foetidum* is a pungently smelling, tropical herb and is used as food, flavouring agent and ethnomedicine [35].









The Temiar communities (i.e. Temiar villages at Pos Brooke) are heavily dependent on rice and are consuming more processed food compared to their ancestors. Additionally, villages situated near to the the main road (Jalan Cameron Highlands-Gua Musang) commonly are able to purchase food items. However, during the flood which wiped out some of the villages connecting bridges and roads, the Temiar communities were affected as food aid were slow to reach their villages. During the flood, the villagers from Kg. Jedip, Pos Brook and Kg. Teranek, Kuala Betis sought for shelter and safety in the nearby hills. Food aid were slow to reach the villages as access to the villagers were inundated by floodwaters. Meanwhile, there were shortage of processed food in Kg. Gob, Pos Gob during the flood. Kg. Gob is situated in the interior of Kelantan jungle bordering with Perak state. Roads were too treacherous (normal dry weather will take 5 hours of travelling in a four wheel drive) and connecting bridges and roads were damaged during the flood as well.







The villagers resorted to harvesting wild, indigenous plants for consumption. In total 14 types of plants and tubers were utilised heavily during the flood and before the villagers were evacuated to relief centres. The head of tribesmen and Tok Batins assisted in identifying the edible plants (i.e. monggoi, betek, kelap), tuber (ubi) and source of clean drinking water (i.e. water from cewes). However, the villagers did not hunt for wild animals during the flood. Based on the in-depth interview, the respondents stated that there were very few sightings of wild animals and they prefer to conserve their energy whilst waiting for evacuation. Additionally, most of the villagers left their blowpipes when seeking higher ground.

During the peak season of the flood (end of December 2014), the villagers depended on the edible, wild plants and tubers to survive. Most of the plants, tubers and rice provided energy and micronutrients for the the villagers. Hence by utilising their local knowledge, the villagers were able to identify wild, edible plants during the flood situation. These indigenous knowledge that are communicated verbally through generations are important. In fact, the villagers ensure their survivability during the flood by correctly identifying edible food

sources for themselves. Indigenous knowledge such as these should be documented to prevent the loss of valuable information.

Table 2: Indigenous plants as alternative food sources during flood season.

| Local Name | Species Name | Parts Used | Preparation Method | Photo | Altitude | Latitude and Longitude |
|--|---------------------------------|--|---|--|----------|----------------------------------|
| <i>Bergentok</i> (Temiar); <i>Kulatelinga</i> (Malay); Ear fungus (English) | <i>Auriculariasp.</i> | Whole fruiting body (note only the brown fungi were identified as alternative food source by the locals) | Harvested and eaten raw. |  | 866 m | 04° 39'01.7" N 101°29'43.8" E |
| <i>Monggoi</i> (Temiar); <i>Terung pipit</i> (Malay); Devil's fig, turkey berry, wild eggplant (English) | <i>Solanumtorvum</i> | Fruits | Fruits are harvested, sliced and boiled / cooked. |  | 729 m | 04° 39.688'N 101°29.274'E |
| <i>Geraloh</i> (Temiar); <i>Berangan</i> (Malay) | <i>Castanopsismegacarpa</i> | Fruits | Slice open cupule to reveal brownish nut. Nuts can be boiled / roasted. |  | 922 m | 04°39'01.7"N 101°29'43.8"E |
| <i>KeberMoi</i> (Temiar); Berries (English) | <i>Rubussumatranus</i> | Fruits | Mature fruits are harvested and eaten raw. |  | 895 m | 04°39'01.7"N 101°29'45.3"E |
| <i>Karyiel; Catak</i> (Temiar) | <i>Molineriacapitulata</i> | Fruits | Fruits are harvested, sliced and boiled / cooked |  | 702 m | 04°39'.688"N 101°29'.274"E |
| <i>Peng Hong</i> (Temiar) | <i>Erechtitesvalerianifolia</i> | Leaf | Young leaves are harvested and cooked / boiled. |  | 828 m | 04°38.920'N 101°29.881'E |
| <i>Cerwes</i> (Temiar); Sappanwood (English) | <i>Caesalpiniasappan</i> | Stem | Branch / stem is chopped and allow water to drip into container. The water extract can be used as alternative drinking water. |  | 749m | 04°39.532'N 101°29.573'E |
| <i>Rebungbetek</i> (Temiar) | <i>Dendrocalamusasper</i> | Shoots | Shoots are cleaned, sliced into small pieces and boiled/stir-fried. |  | 761 m | 04°39.497'N 101°29.572'E |

| | | | | | | |
|--|----------------------------------|---------------|--|--|-------|-------------------------------|
| <i>Cengkarak</i> (Temiar); Koster's curse, clidemia, soap bush (English) | <i>Clidemiahirta</i> | Berries | Fruits are plucked and consumed directly. |  | 891 m | 04°38'.664"N 101°30'.029"E |
| Kuala Betis | | | | | | |
| <i>Sayurkelap</i> (Temiar); <i>Salangsuar</i> (Malay) | <i>Pentaphragmabegoniifolium</i> | Leaves | Leaves are trimmed, washed and cooked. |  | 231 | 04°53'58.6"N 101°46'45.9"E |
| <i>Te-ok</i> (Temiar) | <i>Arengaobtusifolia</i> | Shoots | Stem is sliced to reveal inner whitish, soft shoot. Shoot should be cooked before consumption. |  | 231 | 04°53'58.6"N 101°46'45.9"E |
| Pos Gob | | | | | | |
| <i>Ubikayu</i> (Temiar; Malay) | <i>Manihotesculenta</i> | Cassava roots | Cassava skin is peeled, washed and roasted / boiled. |  | 490 | 05°44'45.6"N 101°51'52.2"E |
| <i>Padihuma / bukit</i> (Temiar) <i>Padibukit</i> (Malay); Upland rice (English) | <i>Oryzaglaberrima</i> | Grains | Paddy is pound to remove husk, washed and boiled. |  | 485 | 05°44'45.6"N 101°51'52.2"E |
| <i>Mong</i> (Temiar) <i>Pokokpijat</i> (Malay); Mexican coriander, spirit weed (English) | <i>Eryngiumfoetidum</i> | Leaves | Leaves are harvested, mashed and stir-fried. |  | 502 | 05°16'07.6"N 101°39'10.9"E |

Conclusion:

The recent flood had forced the local Temiar communities to search for alternative food sources. A total of 14 types of plants and tuber were utilised by the villagers who took shelter in the jungle/hills. With guidance from the head of tribesman, medicine man/woman and elderly communities, the villagers had correctly identified edible wild plants for consumption. Wild local plants were readily available and contributed nutrients and energy to the villagers. This benefit the younger generation of Temiar communities in identifying wild, edible plants that can be utilised by the villages and contribute to the food and nutritional security of the local Temiar community. Nutritional properties of the plants could reveal potential usefulness of these plants.

ACKNOWLEDGEMENT

The authors gratefully acknowledge the FRGS Flood Disaster Management Grant (R/FRGS/A07.00/00295A/003/2015/000251), JAKOA and the Temiar communities of Lojing, Kuala Betis and Pos Gob.

REFERENCES

- [1] The Star., 2015. Collapsed bridges and roads cut aid to villagers.
- [2] Mohammad, B.U., A.M. Sharif, 2012. Ethnomedicinal knowledge of Khasia tribe in Sylhet region, Bangladesh. Indian Journal of Tropical Biodiversity, 20(1): 69-76.
- [3] Odhav, B., S.Beekrum, U.Akula and H. Baijnath, 2007. Preliminary assessment of nutritional value of traditional leafy vegetables in KwaZulu-Natal, South Africa. Journal of Food Composition and Analysis, 20(5): 430-435.
- [4] Gupta, S. and K.Bains, 2006. Traditional cooked vegetable dishes as important sources of ascorbic acid and β -carotene in the diets of Indian urban and rural families. Food and Nutrition Bulletin, 27: 306-310.
- [5] Dias, M.I., L. Barros, R.C. Alves., M.B.P.P. Oliveira, C. Santos-Buelga and I.C.F.R. Ferreira, 2014. Nutritional composition, antioxidant activity and phenolic compounds of wild *Taraxacum*sect.Ruderalia. Food Research International, 56: 266-271.
- [6] Mutheeswaran, S., P.Pandikumar, M.Chellappandian and S.Ignacimuthu, 2011. Documentation and quantitative analysis of the local knowledge on medicinal plants among traditional Siddha healers in Virudhunagar district of Tamil Nadu, India. Journal of Ethnopharmacology, 137: 523-533.
- [7] Toyang, N.J. and R.Verpoorte, 2013. A review of the medicinal potentials of plants of the genus *Vernonia* (Asteraceae). Journal of Ethnopharmacology, 146(13): 681-723.
- [8] Wang, X.M., J.Zhang,L.H. Wu, Y.L. Zhao, T. Li, J.Q. Li, Y.Z. Wang and H.G. Liu, 2014. A mini-review of chemical composition and nutritional value of edible wild-grown mushroom from China. Food Chemistry, 151:279-285.
- [9] Usiku, N.P., A.Oelofse,G.K. Duodu, J.M. Bester and M.Faber, 2010. Nutritional value of leafy vegetables of sub- Saharan Africa and their potential contribution to human health: A review. Critical review in Journal of Food Composition and Analysis, 23: 499-509.
- [10] Schonfeldt, H.C. and B. Pretorius, 2011. The nutrient content of five traditional South African dark green leafy vegetables – A preliminary study. Journal of Food Composition and Analysis, 24(8): 1141-1146.
- [11] Charles, A.L., K.Sriroth and T.C. Huang, 2005. Proximate composition, mineral contents, hydrogen cyanide and phytic acid of 5 cassava genotypes. Food Chemistry, 92(4): 615-620.
- [12] Montagnac, J.A., C.R. Davis and S.A.Tanumihardjo, 2009. Nutritional value of cassava for use as a staple food and recent advances for improvement. Comprehensive reviews in Food Science and Food Safety, 8: 181-194.
- [13] FAO, 1990. Roots, tubers, plantains and bananas in human nutrition. FAO UN: Rome.
- [14] JAKOA, 2014. Sub-tribe/ethnic. Available from: <http://www.jakoa.gov.my/en/web/guest/suku-kaum/bangsa> [Accessed 12 February 2014]
- [15] Semenya, S.S. and A. Maroyi, 2012. Medicinal plants used by the Bapedi traditional healers to treat diarrhoea in the Limpopo Province, South Africa. Journal of Ethnopharmacology, 144:395-401.
- [16] Garibay-Orijel, R., J. Cordova, J. Cifuentes, R.Valenzuela, A. Estrada-Torres and A. Kong, 2009. Integrating wild mushrooms use into a model of sustainable management for ingenuous community forests. Forest Ecology and Management, 258(15): 122-131.
- [17] Lee, S.S., Y.S. Chang and M.N.R. Noraswati, 2009. Utilization of macrofungi by some indigenous communities for food and medicine in Peninsular Malaysia. Forest Ecology and Management, 257(10): 2062-2065.
- [18] Jaiswal, B.S., 2012. *Solanum torvum*: A review of its traditional uses, phytochemistry and pharmacology. International Journal of Pharma and Bio Sciences, 3(4): 104-111.
- [19] Yasuda, M., S.Miura, N. Ishii, T. Okuda and Nor Azman Hussein, 2004. Fallen fruits and terrestrial vertebrate frugivores: a case study in a lowland tropical rainforest in Peninsular Malaysia. In Forget, P.-M., J.E Lambert, P.E. Hulme and S.B. Vander Wall. Seed Fate: Predation, Dispersal and Seedling Establishment. Oxford: CABI Publishing.
- [20] Nguanchoo, V., P. Srisanga, S. Swangpol, S. Prathanturug and T. Jenjittikul, 2014. Food plants in Hmong cuisine in Northern Thailand. Thai Journal of Botany, 6(2): 131-145.
- [21] Panyaphu, K., T.V. On, P. Sirisa-ard, P. Srisa-nga, S. ChansaKaow and S. Nathakarnkitkul, 2011. Medicinal plants of the Mien (Yao) in Northern Thailand and their potential value in the primary healthcare of postpartum women. Journal of Ethnopharmacology, 135(2): 226-237.

- [22] Balangcod, T.D. and A.K.Balangcod, 2009. Underutilized plant resources in Tinoc, Ifugao, Cordillera Administrative Region, Luzon Island, Philippines. *Acta Horticulturae*, 806: 647-654.
- [23] Eswani, N., K.Abd Kudus, M. Nazre, A.G. Awang Noor, 2010. Medicinal plant diversity and vegetation analysis of logged over hill forest of TekaiTembeling Forest Reserve, Jerantut, Pahang. *Journal of Agricultural Science*, 2(3): 189-210.
- [24] Kulip, J., 1996. A survey of indigenous plants used for food and medicine by the Kadazandusun ethnic in Tambunan, Sabah, East Malaysia. Paper presented at Borneo Research Council Fourth Biennial Conference of The Borneo Research Council, 10-15 June, Universiti Brunei Darussalam.
- [25] Shrishailappa, B., M.Sudheer and B. Suresh, 2004. *Caesalpiniasappan* – A medicinal and dye yielding plant. *Natural Product Radiance*, 3(2): 75-82.
- [26] Saenjum, C., C.Chaiyasut, S. Kadchumsang, S. Chansakaowand M.Suttajit, 2003. Antioxidant activity and protective effects on DNA damage of *Caesalpiniasappan* L. extract. *Journal of Medicinal Plants Research*, 4: 1594-1600.
- [27] Safitri, R., P.Tarigan, H.J. Freisleben, R.J.Rumampuk and A.Murakami, 2003. Antioxidant activity in vitro of two aromatic compounds from *Caesalpiniasappan* L. *BioFactors*, 19: 71-77.
- [28] Nirmala, C., D.Elangbam and M.L. Sharma, 2007. Changes in nutrient components during ageing of emerging juvenile bamboo shoots. *International Journal of Food Sciences and Nutrition*, 58(8): 612-618.
- [29] Satya, S., L.M.P. Bal, Singhal and S.N. Naik, 2010. Bamboo shoot processing: food quality and safety aspect (a review). *Trends in Food Science and Technology*, 21(4): 181-189.
- [30] Styger, E., J.E.M.Rakotoarimanana, R.Rabevohitra and E.C.M. Fernandes, 1999. Indigenous fruit trees of Madagascar: potential components of agroforestry systems to improve nutrition and restore biological diversity. *Agroforestry Systems*, 46: 289-310.
- [31] Wiart, C., 2012. Medicinal plants of China, Korea, and Japan: Bioresources for Tomorrow's Drugs and Cosmetics. Boca Raton: CRC Press, 352.
- [32] Singh, R.K., R.C.Srivastava, C.B.Pandey and A. Singh, 2015. Tribal institutions and conservation of the bioculturally valuable 'tasat' (*Arengaobtusifolia*) tree in the eastern Himalaya. *Journal of Environmental Planning and Management*, 58(1): 61-90.
- [33] FAOSTAT, 2013. Cassava. Food and Agriculture Organization.
- [34] Rahmat, Z., A.Wagiran, N. MohdNazir, S. MohdArif, S.N.A. Zulkifli, A. AbdSamad, M.S. Shamsir, F.MohdSalleh and M.R. Sarmidi, 2014. Potensipadibukitsebagaialternatifkepadapadisawah. *Jurnal Teknologi*, 70(6): 89-92.
- [35] Paul, J.H.A., C.E. Seaforth and T. Tikasingh, 2011. *Eryngiumfoetidum* L.: A review. *Fitoterapia*, 82(3): 302-308.