

Preliminary study on leech crude extract as an anti-coagulant agent

Zulhisyam Bin Abdul Kari, Ahmad Anwar Ismail and Ibrahim Che Omar

*Faculty of Agro Industry and Natural Resources,
University Malaysia Kelantan, 16100,
Pengkalan Chepa, Kota Bharu, Kelantan.*

Abstract

Hirudin is the generic name for a family of closely related homologous peptides that are all found in the cranial salivary glands of medicinal leech (Wallis, 1996). In 1984, the blood-thinning agent (anticoagulant) hirudin was isolated for the first time from leech salivary glands (Weinnberg, 1994). Today, hirudin is the most specific and active of known thrombin inhibitors not influencing on other peptidases (Hofsteenge and Stone, 1987). *Hirudinea* sp. (Buffalo Leech) was used for the isolation of hirudin for anti-coagulant application. The leech crude extract was prepared by using conventional method and tested to fresh rabbit blood. The time of blood clotting was recorded by using a drop method (Prakasam *et al.*, 2005). Three replicates were applied for each treatment. The results showed that the rabbit blood treated with leech crude extract gave the longest time of blood to clot with average and standard deviation values of 15.33 ± 0.6 minutes when compared to treatment with distilled water and that on blood only (control) with average and standard deviation values of 2.83 ± 0.3 and 0.59 ± 0.01 minutes, respectively. These results revealed that crude extract of leeches contain an anti-coagulant agent to prevent blood from clotting. The study on leech crude extract will be extended to include purification of hirudin for applications as anti-coagulant.

Keywords: Protein hirudin, leech crude, anticoagulant agent, blood-clotting

Introduction

Traditionally, leeches are widely used as a model animal in toxicological, physiological, neurobiological, biochemical, histological and many other studies (Mann, 1962; Flerov and Lapkina, 1976; Lapkina and Flerov, 1979; Sawyer, 1986; Lapkina, 1992; Huguet and Molinas, 1992, 1996; Blackshaw and Nicholls, 1995; Petrauskienė, 2001). The use of leeches for clinical or medicinal purposes (e.g. blood-letting) has occurred since the 5th century BC, with considerable usage in the 19th century, followed by a decline in the early 20th century (Kasperek *et al.*, 2000). However, in more recent years, the medically beneficial usage of leech is once again increasing (Baskova *et al.*, 1983, 1992). For example, *H. medicinalis* is used by plastic surgeons to restore venous circulation in tissue grafts where blood stagnation is a problem (Sawyer, 1986; Rigbi *et al.*, 1987; Roters and Zebe, 1992; Whitaker *et al.*, 2004; Huang *et al.*, 2006b).

Extracts from this species have been shown to have an important thrombolytic effect on experimental thrombosis (Tan and Liu, 2002; Huang *et al.*, 2006a, 2006b; Li *et al.*, 2006). There has been increasing collection of this species for medicinal purposes in the 20th century, and this, combined with a general loss and pollution of wetland habitats have caused a dramatic decline of *H. manillensis* throughout its geographical range (Steiner *et al.*, 1990; Electricwala *et al.*, 1993; Singhal and Davies, 1996).

In this country, it is not known or proven conclusively that the locally named Buffalo Leech is not of *H. manillensis*. Local taxonomists have not been able to identify the species used those for medical purposes and would rather refer to its genus only as *Hirudinea* sp.

Material and Methods

Sample Preparation

Hirudinea sp. used in the study was provided by PT Dynamic Consultant Co., Kota Bharu, Kelantan. The non-chlorine freshwater in the aquarium tank placed indoor was aerated, with 50% of the water changed once every 3 days. The temperature, pH and light intensity were maintained at room temperature (25° C -27 °C), 6-8, and 0 - 100 lux, respectively. The leeches were starved before starting this study.

Experimental Method

For the experiment, 10 gram of leeches was taken and the crop emptied of the blood by placing the leeches in a container filled with crystals of sodium chloride for 15 minute and then washed with distilled water. They are then cut into small pieces and grinded using a blender. The extraction was made with a volume of distilled water equal to six times the weight of the pulp. After centrifugation at 1500 rpm for 60 minutes at 4 °C, the supernatant fluid was filtered through coarse paper. Finally, the crude extract was stored at 4°C for 24 hours.

Precipitation

A 0.5 % copper sulfate solution was added drop by drop to the prepared extract until the reagent no longer produced further precipitation. After removal of the precipitate by filtration on course paper, the clear solution was placed in visking tubing and dialysed.

Dialysis

Dialysis was done by putting 20 ml sample of the leech extract supernatant into visking tubing and dialysed against cold distilled water at 4°C on the shaking machine for eight hours at 100 rpm. The leech extract was stored at 4 °C for 24 hours.

Testing anti-coagulant power of leech extract on blood

The crude extract was tested on their anti-coagulant power activity by adding 0.5 ml of the extract to 2 ml of fresh rabbit blood and the time of clotting recorded.

Results

Figure 1 show the time for blood to clot for different treatments. Analysis carried out using the One-Way ANOVA showed significant differences among the different treatments.




Figure 1: Blood clotting time under different treatments.

Treatment	Time of blood clotting (minutes)
(1) Rabbit blood only (Control)	0.59±0.01
(2) Rabbit blood + distilled water	2.83±0.3
(3) Rabbit blood + crude extract	15.33±0.6

Table 1: Time taken for blood to clot under three treatment regimes. Data in the table were means and standard deviations (mean±S.D.)

Table 1 shows treatment 3 gave the longest time for blood to clot with mean and standard deviation values of 15.33 ± 0.6 when compared with treatments 1 and 2 with mean and standard deviation values of 0.59 ± 0.01 and 2.83 ± 0.3 , respectively.

Discussion

This experiment was conducted to determine the anti-coagulant activity of in the buffalo leech extract. The anti-coagulant power was evaluated by testing the crude extract on fresh rabbit blood and recording the time of clotting. The fact that a strong anti-coagulant activity was exhibited and the local buffalo leeches indicates that hirudin was present in the crude extract. Generally, Hirudin is the generic name for a family of closely related homologous peptides that are all found in the cranial salivary glands of medicinal leech (Wallis, 1996). In 1984, the blood-thinning agent (anticoagulant) hirudin was isolated for the first time from leech salivary glands (Weinnberg, 1994). Today, hirudin is the most specific and active of known thrombin inhibitors not influencing on other peptidases (Hofsteenge and Stone, 1987).

Figure 1 represents the time of blood to clot at different treatments showed very significantly different ($p=0.00$). Treatment 3 gave the highest result with mean and standard deviation values of 15.33 ± 0.6 compared with treatment 1 gave the lowest result with mean and standard deviation values of 0.59 ± 0.01 .

In this study they was marked differences in the clotting time for the treatments employed with that treated on rabbit blood gave the highest values. The result of this preliminary study revealed that crude extract of local leeches contained an anti-coagulant to prevent blood from clotting.

Acknowledgements

The author is grateful to the Faculty of Agro Industry and Natural Resources for providing space and facilities to carry out the study. This study was funded by the University Malaysia Kelantan and supported by PT Dynamic Consultant Co., Kota Bharu, Kelantan and Freshwater Fisheries Research Centre, Gelami Lemi, Jelebu, Negeri Sembilan. For the statistical analysis my grateful thanks are extended to En. Che Jaffar Sulong for advice and help.

References

- Baskova, I.P., Khalil, S., Nartikova, V.F., Paskhina, T.S., 1992. Inhibition of plasma kallikrein. kininase and kinin-like activities of preparations from the medicinal leeches. *Thrombosis Research* 67 (6), 721–730.

- Baskova, I.P., Cherkesova, D.U., Mosolov, V.V., 1983. Hirudin from leech heads and whole leeches and “pseudo-hirudin” from leech bodies. *Thrombosis Research* 30 (5), 459–467.
- Blackshaw, S.E., Nicholls, J.G., 1995. Neurobiology and development of the leech. *Journal of Neurobiology* 27 (3), 267–276.
- Electricwala, A., Hartwell, R., Scawen, M.D., Atkinson, T., 1993. The complete amino acid sequence of a hirudin variant from the leech *Hirudinea manillensis*. *Journal of Protein Chemistry* 12 (3), 365-370.
- Flerov, B.A., Lapkina, L.N., 1976. Avoidance of some toxicants by medicinal leech. *Inform. Bull. IBVV*. 30, 48–52 (In Russian).
- Huang, A.M., Li, Z.Y., Liao, G.S., Huang, H.K., Ban, J.D., Lin, F.Q., 2006b. The anticoagulant effect of Bufrudin on human plasma in vitro and rabbit plasma in vivo. *Journal of Guangxi Medical University* 23 (1), 30–32 (In Chinese).
- Huang, A.M., Li, Z.Y., Liao, G.S., Ban, J.D., 2006a. Purification of anti-coagulant protein from the digestive juice of Guangxi *Hirudinaria manillensis*. *Chinese Journal of Biochemical Pharmaceutics* 27 (5), 273-256 (In Chinese).
- Huang, A.M., Li, Z.Y., Liao, G.S., Huang, H.K., Ban, J.D., Lin, F.Q., 2006b. The anticoagulant effect of Bufrudin on human plasma in vitro and rabbit plasma in vivo. *Journal of Guangxi Medical university* 23 (1), 30-32 (In Chinese).
- Huguet, G., Molinas, M., 1992. Changes in epithelial cells in *Hirudo medicinalis* during wound healing. *Journal of Invertebrate Pathology* 59, 11–17.
- Huguet, G., Molinas, M., 1996. Myofibroblast-like cells and wound contraction in leech wound healing. *Journal of Experimental Zoology* 275, 308–316.
- J. Hofsteenge, S.R. Stone, The effect of thrombomodulin on the cleavage of Wbrinogen and Wbrinogen fragments by thrombin, *Eur. J. Biochem.* 168 (1987) 49–56.
- Kasperek, M., Demirsoy, A., Akbulut, A., Akbulut, N.E., Çalışkan, M., Durmuş, Y., 2000. Distribution and status of the medicinal leech (*Hirudo medicinalis* L.) in Turkey. *Hydrobiologia* 441, 37–44.
- Lapkina, L.N., 1992. Comparative study of lethal and sublethal effects of trichlorphon on leeches. *Information bulletin IBVV. RAN.* 94, 67-73 (In Russian).
- Lapkina, L.N., Flerov, B.A., 1979. Impact of some toxicants on leech in acute toxicity test. *Proc. Freshwater Biol. Inst. Academic Science USSR.* 38 (41), 50–59 (In Russian).
- Li, Y.H., Yang, Li, Z.Y, 2006. The antithrombotic effect of extract from the Guangxi *Hirudinaria manillensis* on animal thrombosis. *Journal of Guangxi Medical University* 23 (6), 901-903 (In Chinese).

- Mann, K. H. 1962. *Leeches (Hirudinea): Their structure, physiology, ecology and embryology*. The Macmillan Company, New York, USA.
- Petrauskienė, L., 2001. Water toxicity assessment using medicinal leeches. *Aquatic Ecosystem Health & Management* 4, 203–208.
- Rigbi, M., Levy, H., Iraqi, F., Teitelbaum, M., Orevi, M., Alajoutsijärvi, A., Horovitz, A., Galun, R., 1987. The saliva of the medicinal leech *Hirudo medicinalis* I. Biochemical characterization of the high molecular weight fraction. *Comparative Biochemistry and Physiology B.*, 87 (3), 567–573.
- Roters, F.J., Zebe, E., 1992. Proteinases of the medicinal leech, *Hirudo medicinalis*: purification and partial characterization of three enzymes from the digestive tract. *Comparative Biochemistry and Physiology B.*, 102,627–634.
- S.L. Weinberg, New medicine from old-hirudin and the leech, *J. Am. Coll. Cardiol.*, 23 (1994) 544.
- Sawyer, R. T. 1986. *Leech biology and behaviour*. Volumes I, II, III. Clarendon Press, Oxford, England.
- Singhal, R.N., Daviers, R.W, 1996. Effect of an organophosphorus insecticide (Temephos) on gametogenesis in the leech *Hirudinaria manillensis* (Hirudinidae). *Journal of Invertebrate Pathology*, 67 (1), 100-101.
- Steiner, V., Knecht, R., Gruetter, M., Raschdorf, F., Gassmann, E., Maschler, R., 1990. Isolation and purification of novel hirudins from the leeches *Hirudinaria manillensis* by high-performance liquid chromatography. *Journal of Chromatography B: Biomedical Sciences and Application* 530, 273-283.
- Tan, E.G., Liu, X.P., 2002. Cloning and sequencing of Hirudin gene of *Hirudinaria manillensis* in Guangdong, China. *Academic Journal SUMS* 23 (2) 84-86 (In Chinese)
- Wallis RB. Hirudin: From leeches to man. *Sem Thromb Hemost*, 1996; 22: 185–196.
- Whitaker, I.S., Izadi, D., Oliver, D.W., Monteath, G., Butler, P.E., 2004. *Hirudo medicinalis* and the plastic surgeon. *British Journal of Plastic Surgery*, 57, 348–353.